



A CHANCE TO LOWER THE RISK OF CANCER SPREADING OR RETURNING

Actor portrayals. OPDIVO is not approved for patients younger than 18 years of age.



Given before surgery to adults with non-small cell lung cancer to help prevent it from returning after surgery

OPDIVO® will not work for everyone. Individual results may vary. Please see information about the clinical trial, including serious side effects, on pages [4-5](#).

What is OPDIVO?

OPDIVO is a prescription medicine used in combination with chemotherapy that contains platinum and another chemotherapy medicine, before you have surgery, for adults with **early-stage lung cancer (called non-small cell lung cancer)**.

OPDIVO is not approved for patients younger than 18 years of age.

Information provided in this brochure is not a substitute for talking with your healthcare professional. Your healthcare professional is the best source of information about your disease.

OPDIVO[®]
(nivolumab)

INJECTION FOR INTRAVENOUS USE 10 mg/mL



Please see Important Facts about OPDIVO, including serious side effects, on pages [7-8](#) and [U.S. Full Prescribing Information and Medication Guide](#).

First + Only immunotherapy-based option before surgery for early-stage non-small cell lung cancer

OPDIVO is an immunotherapy. That means it works with your own immune system to fight cancer.



This immunotherapy is made by Bristol Myers Squibb, which has been at the forefront of the battle against cancer since 1967, recently paving the way in the field of immuno-oncology.

Today, OPDIVO + chemo could enhance your treatment plan with the first-and-only FDA-approved immunotherapy combination given before surgery for early-stage non-small cell lung cancer.

It's also yet another milestone in our continued commitment to help people living with this condition.


**Ask your doctor today if OPDIVO + chemo
before surgery is right for you**

i Please see additional Important Facts about OPDIVO, including serious side effects, on pages 7-8 of the brochure and [U.S. Full Prescribing Information and Medication Guide](#).


Help Seek and Attack

How OPDIVO works with your immune system

OPDIVO
(nivolumab)
INJECTION FOR INTRAVENOUS USE 10 mg/mL



OPDIVO can cause your T cells to attack healthy cells contained within organs and tissues in any area of your body and can affect the way they work. These problems can sometimes become serious or life-threatening and can lead to death, and may happen anytime during treatment or even after treatment has ended. You may have more than one of these problems at the same time. Your doctor will check you for these problems during treatment.

 Visit [OPDIVO.com](https://www.opdivo.com) for more information.

 Please see Important Facts about OPDIVO, including serious side effects, on pages 7-8 and [U.S. Full Prescribing Information and Medication Guide](#).

A Head Start – Treatment before surgery gives you and your doctor an extra option

What is OPDIVO?

OPDIVO is an immunotherapy.

What is immunotherapy?

Immunotherapy is a treatment that works with your immune system to fight cancer.

How does OPDIVO treat early-stage non-small cell lung cancer?

OPDIVO is combined with chemotherapy to treat this cancer before surgery.

Why consider OPDIVO + chemo?

Treatment with OPDIVO + chemo before surgery could offer you an additional way to treat your condition because it attacks cancer cells that might remain "hidden" or hard to see.

PRE-SURGERY OPTIONS

To make sure the tumor is small enough to be removed, your doctor may recommend a pre-surgery treatment.

OPDIVO + CHEMO

or

CHEMO and/or RADIATION

SURGERY

Your oncologist shares the results of your treatments

CHEMOTHERAPY: This treatment uses drugs to stop the growth of cancer cells

RADIATION: This treatment uses high-energy rays and depends on the stage and size of tumors

Ask your oncologist about adding OPDIVO + chemo to your treatment plan

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

- lung problems
- intestinal problems
- liver problems
- hormone gland problems
- kidney problems
- skin problems
- problems in other organs and tissues
- severe infusion reactions
- complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic)

A Chance for Successful Surgery


OPDIVO + chemo delivered stronger results than chemo alone

In a clinical trial of **358** patients with early-stage non-small cell lung cancer who were able to have surgery, **179** patients were given OPDIVO and chemotherapy before surgery while **179** were given chemotherapy.

Patients on OPDIVO + chemo versus chemo alone



OPDIVO + chemo, given before surgery, is proven in a clinical trial to reduce the risk of cancer spreading or returning by 37% compared to chemo alone.



At 31.6 months
half of the patients on OPDIVO + chemo remained free of cancer spreading or returning

At 20.8 months
half of the patients on chemo alone remained free of cancer spreading or returning

10X more patients who were given OPDIVO + chemo before surgery saw tumors disappear completely than those on chemo alone after the surgery.

24% vs **2.2%**
of patients on OPDIVO + chemo vs of patients on chemo alone

OPDIVO will not work for everyone. Individual results may vary.

What are the most common side effects of OPDIVO?


The most common side effects of OPDIVO when used in combination with chemotherapy include:

- nausea
- constipation
- feeling tired
- decreased appetite
- rash

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

Treatment Planning Prior to Surgery

A dosing schedule with 3 infusions every 3 weeks



OPDIVO + chemo is given through intravenous (IV) infusion. The medicine is given directly into the bloodstream through a needle placed in a vein by a healthcare professional—usually in the arm or hand.

Talk to your doctor about which treatment schedule may be right for you. Your doctor will decide how many treatments you need. Your healthcare provider will do blood tests to check you for side effects, and may treat you with corticosteroid or hormone replacement medicines, and delay or completely stop treatment if you have side effects that are severe. Be sure to talk to your healthcare team about side effects.

Frequent communication with your healthcare team is an important part of treatment. Use your appointments to let them know how you're feeling and ask any questions you may have.

If you miss an appointment, call a healthcare provider right away. Try to schedule your appointments for the same day of the week and the same time of day to make them easier to remember. It is important for you to keep all appointments with your healthcare provider.


Please see additional Important Facts about OPDIVO, including serious side effects and information about pregnancy, nursing, health problems, and other medicines that you should discuss with your healthcare team before starting treatment, on pages 7-8 and [U.S. Full Prescribing Information and Medication Guide](#).


Important Facts About OPDIVO


This is a summary of important information that you need to know about OPDIVO. Your healthcare team can work with you to help answer any questions you may have about OPDIVO. Keep this information in a safe place so you can refer to it before and during your treatment.



Look out for the following icons as you read:

 Talk to your healthcare team

 Call a healthcare provider right away

 Helpful information to remember


What is OPDIVO?

OPDIVO is a prescription medicine used in combination with chemotherapy that contains platinum and another chemotherapy medicine, before you have surgery, for adults with **early-stage lung cancer (called non-small cell lung cancer)**.

It is not known if OPDIVO is safe and effective in children younger than 12 years of age with MSI-H or dMMR metastatic colorectal cancer, **or** in children younger than 18 years of age for the treatment of any other cancers.

What is the most important information I should know about OPDIVO?

OPDIVO is a medicine that may treat certain cancers by working with your immune system. OPDIVO can cause your immune system to attack normal organs and tissues in any area of your body and can affect the way they work. Some of these problems may happen more often when OPDIVO is used in combination with another therapy.

 **Get medical help immediately** if you develop any of these signs or symptoms or they get worse. It may keep these problems from becoming more serious. Your healthcare team will check you for these problems during treatment and may treat you with corticosteroid or hormone replacement medicines. If you have severe side effects, your healthcare team may also need to delay or completely stop your treatment.

What are the serious side effects of OPDIVO?

A **serious side effect** is a side effect that can sometimes become severe or life-threatening and can lead to death. They may happen anytime during treatment or even after your treatment has ended. You may have more than one of these problems at the same time.

Call or see your healthcare provider right away if you develop any new or worse signs or symptoms, including:

Lung problems – Things to look out for may include:

- new or worsening cough
- shortness of breath
- chest pain

Intestinal problems – Things to look out for may include:

- diarrhea (loose stools) or more frequent bowel movements than usual
- stools that are black, tarry, sticky, or have blood or mucus
- severe stomach-area (abdominal) pain or tenderness

Liver problems – Things to look out for may include:

- yellowing of your skin or the whites of your eyes
- severe nausea or vomiting
- pain on the right side of your stomach area (abdomen)
- dark urine (tea colored)
- bleeding or bruising more easily than normal

Hormone gland problems – Things to look out for may include:

- headaches that will not go away or unusual headaches
- eye sensitivity to light
- eye problems
- rapid heartbeat
- increased sweating
- extreme tiredness
- weight gain or weight loss
- feeling more hungry or thirsty than usual
- urinating more often than usual
- hair loss
- feeling cold
- constipation
- your voice gets deeper
- dizziness or fainting
- changes in mood or behavior, such as decreased sex drive, irritability, or forgetfulness

Kidney problems – Things to look out for may include:

- decrease in your amount of urine
- blood in your urine
- swelling of your ankles
- loss of appetite

Skin problems – Things to look out for may include:

- rash
- itching
- skin blistering or peeling
- painful sores or ulcers in the mouth or nose, throat, or genital area

Problems can also happen in other organs and tissues. These are not all the signs and symptoms of immune system problems that can happen with OPDIVO. Call or see your healthcare provider right away for any new or worsening signs or symptoms, which may include:

- Chest pain, irregular heartbeat, shortness of breath, swelling of ankles
- Confusion, sleepiness, memory problems, changes in mood or behavior, stiff neck, balance problems, tingling or numbness of the arms or legs
- Double vision, blurry vision, sensitivity to light, eye pain, changes in eyesight
- Persistent or severe muscle pain or weakness, muscle cramps
- Low red blood cells, bruising

Important Facts About OPDIVO (continued)


What are the possible side effects of OPDIVO?

OPDIVO can cause serious side effects, including:

See the previous section, **“What is the most important information I should know about OPDIVO?”**

Severe infusion reactions – Things to look out for may include:

- chills or shaking
- shortness of breath
- feel like passing out
- itching or rash
- or wheezing
- fever
- flushing
- dizziness
- back or neck pain

 **Tell your healthcare team** right away if you get these symptoms during an infusion of OPDIVO.

Complications of bone marrow (stem cell) transplant that uses donor stem cells (allogeneic). These complications can be severe and can lead to death. These complications may happen if you underwent transplantation either before or after being treated with OPDIVO. Your healthcare provider will monitor you for these complications.

What are the most common side effects of OPDIVO?

The most common side effects of OPDIVO when used in combination with chemotherapy include:

- nausea
- feeling tired
- rash
- constipation
- decreased appetite

These are not all the possible side effects. **Talk to your healthcare team** or pharmacist for more information. You are encouraged to report side effects of prescription drugs to the FDA. Call 1-800-FDA-1088.

What should I discuss with my healthcare team before receiving OPDIVO?

 **Tell your healthcare provider about all your medical conditions, including if you:**

- have immune system problems such as Crohn’s disease, ulcerative colitis, or lupus
- have received an organ transplant
- have received or plan to receive a stem cell transplant that uses donor stem cells (allogeneic)

- have received radiation treatment to your chest area in the past and have received other medicines that are like OPDIVO
- have a condition that affects your nervous system, such as myasthenia gravis or Guillain-Barré syndrome
- are pregnant or plan to become pregnant. OPDIVO can harm your unborn baby.
- are breastfeeding or plan to breastfeed. It is not known if OPDIVO passes into your breast milk. Do not breastfeed during treatment with OPDIVO and for 5 months after the last dose of OPDIVO.



Females who are able to become pregnant:

Your healthcare provider should do a pregnancy test before you start receiving OPDIVO.

- You should use an effective method of birth control during your treatment and for at least 5 months after the last dose of OPDIVO. Talk to your healthcare provider about birth control methods that you can use during this time.
- Tell your healthcare provider right away if you become pregnant or think you may be pregnant during treatment with OPDIVO.



Tell your healthcare provider about all the medicines you take, including:

- prescription medicines
- vitamins
- over-the-counter medicines
- herbal supplements



For more information, please see [U.S. Full Prescribing Information and Medication Guide for OPDIVO](#) or talk to your healthcare team.

Financial Resources

Patient Access, Reimbursement, and Co-Pay Support

**Available through
Bristol Myers Squibb Access Support®**

Bristol Myers Squibb is committed to helping patients gain access to their prescribed BMS medications. That's why we offer the BMS Access Support program, which provides resources to help patients understand their insurance coverage and find information on sources of financial support, including co-pay assistance for eligible commercially insured patients.

For more information, ask your doctor, visit [BMS Access Support](#), or call BMS Access Support at **1-800-861-0048**, 8 AM to 8 PM ET, Monday-Friday.

Other Patient Resources*



GO2 Foundation for Lung Cancer

www.go2foundation.org

GO2 Foundation for Lung Cancer transforms survivorship as the world's leading organization dedicated to saving, extending, and improving the lives of those vulnerable, at risk, and diagnosed with lung cancer. We work to change the reality of living with lung cancer by ending stigma, increasing public and private research funding, and ensuring access to care.



Lung Cancer Research Foundation

www.lungcancerresearchfoundation.org | 844-835-4325

A partnership of lung cancer survivors, advocates, researchers, healthcare professionals, and industry leaders united in the belief that every person with lung cancer deserves a cure.



LUNgevity

www.lungevity.org

A nationally leading lung cancer-focused nonprofit dedicated to changing outcomes for people with lung cancer through research, education, and support.



Lung Cancer Foundation of America (LCFA)

www.lcfamerica.org

An organization committed to improving the survivorship of lung cancer patients through the funding of transformative science.

**Bristol Myers Squibb does not endorse these organizations. The information/links provided by Bristol Myers Squibb are meant for informational purposes only and are not meant to replace a physician's medical advice.*



Please see Important Facts about OPDIVO, including serious side effects, on pages 7-8 and [U.S. Full Prescribing Information and Medication Guide](#).

SIGN UP for OPDIVO with You.

OPDIVO with You means just that: we're here to help you better understand OPDIVO, whether you or a loved one are just considering OPDIVO or already have an OPDIVO prescription. This complimentary program includes the following resources. Sign up and you'll receive:

Helpful Information

Get useful tips, tools, and educational materials that can help you learn more about OPDIVO and may help support your care.

Support and Advocacy Groups

We can direct you to organizations that may be helpful for the challenges you may face.

A Care Counselor as Close as Your Phone

If you are prescribed OPDIVO and sign up for this additional resource, you'll be able to speak with a registered nurse. This personal Care Counselor can answer questions about treatment and help you to be a more informed partner with your doctor. Care Counselors cannot provide medical advice. Your healthcare professional is the best source of information about your health.

If you'd like to sign up for the **OPDIVO with You** program, or if you have general questions about OPDIVO, call **1-855-OPDIVO-1** (1-855-673-4861) 8 AM to 8 PM ET, Monday to Friday, or visit www.OPDIVOwithYou.com



For more information,
call **1-855-OPDIVO-1** or visit
www.OPDIVO.com



SHARE TO INSPIRE

Because your story can help inspire another.

Call us toll-free at **1-855-436-5866**

 Please see Important Facts about OPDIVO, including serious side effects, on pages 7-8 and [U.S. Full Prescribing Information](#) and [Medication Guide](#).

Bristol Myers Squibb is committed to helping patients throughout their treatment.



© 2022 Bristol-Myers Squibb Company.
OPDIVO®, Access Support, and the related logos
are trademarks of Bristol-Myers Squibb Company.

1506-US-2200121 04/22



OPDIVO[®]
(nivolumab)

INJECTION FOR INTRAVENOUS USE 10 mg/mL